

STRATIGRAPHY, PALEONTOLOGY, AND ISOTOPIC AGES OF UPPER MESOZOIC ROCKS IN THE SOUTHWESTERN WRANGELL MOUNTAINS, ALASKA

By ARTHUR GRANTZ, D. L. JONES, and M. A. LANPHERE,
Menlo Park, Calif.

Abstract.—Reconnaissance field observations, fossils, and potassium-argon ages have provided new information on the upper Mesozoic strata of the Wrangell Mountains. The **Kotsina** Conglomerate is probably **Middle** or lower Upper Jurassic. Sandstone along the Chetaslina River may be **Callovian** (Jurassic), and nearby **unfossiliferous** shale and sandstone is probably also upper Mesozoic. A unit of sandstone, **siltstone**, conglomerate, and calcarenite, previously thought to be **gradational** downward into the **Kotsina** Conglomerate, yielded **Hauterivian** and Barremian (Lower Cretaceous) fossils. Similar rocks occur in the northern Chugach Mountains and at **Kuskulana** Pass, where they rest on granodiorite dated as 141 ± 5 m.y. Sandstone and siltstone with Albian (Lower Cretaceous) fossils rest on the Hauterivian to Barremian rocks at Kuskulana Pass, and shale and arkose of similar age rest on **granodiorite** dated as 126 ± 4 m.y. near Mount Drum.

The southwestern **Wrangell** Mountains, and the adjacent lower Chitina Valley, occupy an important position between geologically better known areas—the upper Chitina Valley to the southeast, and the Nelchina area to the west (fig. 1). A reconnaissance survey of the upper Mesozoic rocks in the report area was made to determine regional stratigraphic relationships and the age of certain poorly dated formations. Knowledge of these relationships may have an economic application in the nearby **Copper River** lowland, where poorly exposed upper Mesozoic rocks are a possible source of petroleum.

The upper Mesozoic rocks rest unconformably upon altered upper Paleozoic sedimentary and volcanic rocks, Triassic basalt, and Upper Triassic limestone, black shale, and argillite. The pre-Jurassic rocks were strongly folded, intruded by plutonic rocks, and uplifted and deeply eroded by late Mesozoic time. They supplied most of the detritus found in the upper Mesozoic formations. The upper Mesozoic formations




FIGURE 1.—Extent of upper Mesozoic rocks of the Matanuska geosyncline (shaded area) and area discussed in text. 1, Nelchina area; 2, Copper River lowland; 3, southwestern Wrangell Mountains and lower Chitina Valley, (report area); and 4, upper Chitina Valley.

occupy a synclinal belt between the Chokosna River and Mount Drum and rest on Paleozoic rocks in a small area of the Chugach Mountain front near Chitina (fig. 2).

PREVIOUS WORK

Pioneer reconnaissance surveys by Rohn (1900), Schmdler and Spencer (1901), and Moffit and Maddren




FIGURE 2—Generalized geologic map of the southwestern Wrangell Mountains, Alaska, and generalized columnar sections of Cretaceous rocks in the area. Map after Moffit (1938). Large letters on map refer to stratigraphic sections on facing page.




FIGURE 2.—Continued

(1909) resulted in the recognition of shale and limestone of supposed Late Jurassic or Early Cretaceous age in the Kotsina-Kuskulana Rivers area. These rocks were considered to be part of the Kennicott Formation, and of the thick Kotsina Conglomerate (Rohn, 1900, p. 431) of unknown age. In addition, Mendenhall (1905) mapped unaltered sedimentary rocks which he thought mere Tertiary, in the Cheshnina, Chetaslina, and Dadina River drainage areas, but these rocks are now known to be upper Mesozoic. Near the mouth of the Chitina River, Moffit (1914) found two units of supposed Middle Jurassic age, a unit of "tuffaceous slate" overlying a "tuffaceous conglomerate".

In the Kotsina-Kuskulana Rivers district, Moffit and Mertie (1923) recognized three units of upper Mesozoic rocks to which they assigned a Late Jurassic age. The lowest unit is the Kotsina Conglomerate which unconformably overlies Upper Triassic shale and consists of massive conglomerate with rare shale lenses. The Kotsina was thought to be overlain by an unnamed unit of sandstone, conglomerate, and shale, and this in turn by an unnamed light-colored, highly fossiliferous limestone unit. Moffit and Mertie did not retain the name Kennicott Formation for the upper two units.

The sporadic distribution of these three units, and the fact that in no place mere the upper two found in

stratigraphic sequence with the Kotsina Conglomerate, led to uncertainty as to their relationships. The massive Kotsina Conglomerate was thought to grade laterally into soft yellowish-brown fossiliferous sandstone of the middle unit to the north. The lack of thick conglomerate in some places between the upper limestone unit and the Triassic formations, however, suggested that either the Kotsina was deposited in a restricted area, or that it was removed by erosion prior to deposition of the limestone.

Moffit (1938) published a summary of the geology of the entire Chitina Valley and adjacent area, but the relationships of the upper Mesozoic rocks of the Kotsina-Kuskulana Rivers area and nearby areas to those in other parts of the region were puzzling; they could not be correlated either lithologically or faunally with any other known sequence. Likewise, the supposed Middle Jurassic tuffaceous beds exposed near the mouth of the Chitina River could not be correlated with any others within the Chitina Valley region.

Imlay and Reeside (1954, p. 231) correlated the Kotsina Conglomerate with the basal part of the Matanuska Formation of supposed Coniacian (Late Cretaceous) age. The fossiliferous limestone was not mentioned and presumably was also regarded as part of the Matanuska Formation. They also assigned an Albian (Late Cretaceous) age to the Kennicott Formation, which crops out eastward from Kuskulana Pass, at the east margin of the present study area.

PRESENT INVESTIGATIONS

This report is based on 9 days of helicopter-supported fieldwork carried out by Grantz in the summer of 1963, and contains the results of paleontologic studies by Jones and isotopic age measurements by Lanphere. An abstract of the data resulting from this collaborative work has been published previously (Grantz and others, 1965). Geologic observations were made throughout an area of about 1,200 square miles, but time did not permit tracing contacts or thorough examination of all exposures.

It is assumed that the potassium-argon ages discussed are at least minimum ages, because there is no evidence that biotite or hornblende from granitic rocks incorporate significant amounts of radiogenic argon at the time of crystallization. The plus-or-minus value assigned to each age (table 1) is the estimated standard deviation of analytical precision.

Where possible the formations discussed below will be correlated with the standard stages of northwestern Europe; a list of these stages for the Upper Jurassic and Cretaceous is presented in table 2.

JURASSIC ROCKS

Kotsina Conglomerate

The Kotsina Conglomerate, named by Rohn (1900, p. 431) for exposures on and near the Kotsina River, also crops out near the headwaters of Strelna Creek,

TABLE 1.—Potassium-argon ages and analytical data
[Potassium analyses by H. C. Whitehead; argon analyses by M. A. Lanphere]

Sample No.	Sample description	Mineral	K20 analyses (percent)	Average K ₂ O (percent)	A _{T_{rad}⁴⁰} (10 ⁻¹⁰ moles/g)	$\frac{A_{T_{rad}^{40}}}{A_{T_{total}^{40}}}$	Apparent age (millions of years)	Field No.	Location
1----	Biotite granodiorite clast in Kotsina Conglomerate.	Biotite----	3.97, 3.97	3.97	9.633	0.90	157 A 6	63ALe 14b	Valdez (C-1) quadrangle; lat 61°44'00'' N., long 144°02'50'' W. (loc. H, fig. 2; fig. 3).
2----	Porphyritic hornblende microdiorite dike cutting Kotsina Conglomerate.	Hornblende.	.668, .672	.670	1.460	.83	142 ± 5	63AGz 213	Valdez (C-1) quadrangle; lat 61°44'03'' N., long 144°03'02'' W. (loc. H, fig. 2; fig. 3).
3----	Granodiorite near Kuskulana Pass.	do-----	.767, .784	.776	1.684	.87	141 ± 5	63ALe 15	McCarthy (C-8) quadrangle; lat 61°32'45'' N., long 143°42'30'' W. (loc. E, fig. 2).
4----	Biotite granodiorite near Mount Drum.	Biotite----	7.29, 7.30	7.30	14.11	.92	126 ± 4	63AGz 162	Gulkana (A-2) quadrangle; lat 62°03'09'' N., long 144°34'40'' W. (loc. A, fig. 2).

Decay constants for K⁴⁰: $\lambda_{\gamma} = 0.585 \times 10^{-10}$ year⁻¹; $\lambda_{\beta} = 4.72 \times 10^{-10}$ year⁻¹.
Atomic abundance of K⁴⁰ = 1.19×10^{-4} .

TABLE 2.—Commonly used stage names of the Upper Jurassic and Cretaceous, including the stage names used in this report¹

Series	Stage	
Upper Cretaceous	Maestrichtian	
	Campanian	
	Santonian	
	Coniacian	
	Turonian	
	Cenomanian	
Lower Cretaceous	Albian	
	Aptian	
	Neocomian	Barremian
		Hauterivian
		Valanginian
		Berriasian
Upper Jurassic	Portlandian	
	Kimmeridgian	
	Oxfordian	
	Callovian	

¹ Compiled from Imlay (1952, pl. 2) and Imlay and Reeside (1954, pl. 1).

and was found on the north side of the Cheshnina River valley during the present study (fig. 2).

The physiographic expression of the formation was aptly described by Moffit and Mertie (1923, p. 45) who noted that "The conglomerate mountains are rugged, with precipitous cliffs and a ragged skyline. Their dark color and rough surface give them a forbidding aspect, and in fact many of the ridges are practically impassable." The Kotsina is a very thick bedded, well-rounded, pebble-and-cobble conglomerate that contains boulders in some beds. Hand specimens are prevailing dark or olive gray. The matrix is lithic or feldspathic sandstone which is poorly to fairly well sorted, or in some places dark, hard lutite. The formation contains interbeds and lenses of similar sandstone and of black lutite that commonly is carbonaceous and contains plant scraps. Some of the interbeds are tens of feet thick. The conglomerate clasts are principally dark rocks and of local origin, mostly altered volcanic rocks and Triassic limestone, argillite, and chert. Some light-colored and some mafic plutonic rocks, sandstone, and a little quartz are also present. The size of the clasts appears to decrease toward the southeast. The Kotsina Conglomerate rests unconformably upon rocks as young as latest Triassic. It is probably at least 2,000 to 2,500

feet thick, but its upper contact has not been recognized, and its full thickness is not known. Its estimated thickness and coarseness of grain size suggest that the Kotsina Conglomerate was deposited in a fault- or flexure-bounded basin, probably on or near shore.

The only fossils found in the Kotsina Conglomerate are plant scraps in the lutite layers and Late Triassic mollusks in limestone clasts. Earlier workers based its age upon mollusks found in supposedly equivalent or gradationally overlying marine clastic rocks now known to be of late Neocomian Early Cretaceous (Hauterivian) age. These fossiliferous rocks are unrelated to the Kotsina Conglomerate, and in fact could not be found in contact with it. The supposedly equivalent rocks have been variously considered to be Late Jurassic, Jurassic or Cretaceous, Early Cretaceous, or Late Cretaceous, and the age of the Kotsina Conglomerate varied with the age assigned to these beds.

The Kotsina Conglomerate is intruded by porphyritic hornblende microdiorite dikes that cut a post-Kotsina fault north of the Kotsina River (locality H, fig. 2) and are themselves unconformably overlain by beds of Hauterivian age. These relations are shown on figure 3. Assuming that a significant period of time was required to fault the Kotsina Conglomerate, intrude the dikes, erode the rocks on the west side of the fault to the level of the dikes, and then to submerge the area to receive the Hauterivian sediments, then a Jurassic age for the Kotsina Conglomerate seems likely.

Support for, and considerable refinement of, the stratigraphic age of the Kotsina Conglomerate is




FIGURE 3.—Schematic geologic section showing the field relations critical to dating the Kotsina Conglomerate. Ages were determined by the potassium-argon method. The section is located on the north side of the Kotsina River (loc. H, fig. 2). Geologic units are identified by letter symbol on figure 2.

provided by potassium-argon ages determined for hornblende in one of the microdiorite dikes (sample 2, table 1) which cut the conglomerate and in a biotite granodiorite clast (sample 1, table 1) from the conglomerate in the same outcrop. The age determined for hornblende from the dike is 142 ± 5 million years, and that for biotite from the clast is 157 ± 6 m.y. These determinations indicate a minimum age of Late Jurassic and suggest a Middle or Late Jurassic age for the Kotsina Conglomerate. The biotite age is in good agreement with potassium-argon ages of biotite and hornblende from plutons in the Talkeetna Mountains (Grantz and others, 1963) and the Aleutian Range (Detterman and others, 1965). The stratigraphic evidence in these two areas suggests that emplacement of these plutons occurred in latest Early to earliest Middle Jurassic time. The biotite age for the Kotsina Conglomerate clast suggests that the plutonism which was widespread in the area of the Matanuska geosyncline also occurred locally in the area of the Kotsina Conglomerate.

An age of 141 ± 5 m.y. (sample 3, table 1) was determined for hornblende from a granodiorite pluton near Kuskulana Pass (*loc. E, fig. 2*). This pluton intrudes latest Triassic rocks and is overlain unconformably by Lower Cretaceous (Hauterivian) rocks. Its indicated age is significantly less than the ages that have been measured for granitic rocks in the Talkeetna Mountains and Aleutian Range, which were noted above.

The ages of the hornblende from the pluton at Kuskulana Pass and from the dike which intrudes the Kotsina Conglomerate north of the Kotsina River (samples 2 and 3, table 1) are, however, in excellent agreement, and are considered to represent a Late Jurassic plutonic episode.

Sandstone and lutite

Other rocks in the southwestern Wrangell Mountains that may be Jurassic crop out in the bluffs along the Chetaslina and Cheshnina Rivers (*fig. 2*). These rocks consist mostly of sandstone and lutite and dip 25° to 30° SW. The outcrops along the Cheshnina River lie down-dip from nearby outcrops of the Kotsina Conglomerate which they resemble more closely in degree of induration and deformation than they do the Cretaceous rocks of the area.

The most abundant rock type is greenish-gray, thick-bedded feldspathic and lithic graywacke that is locally cross-bedded and calcareous. It ranges from fine-grained to very coarse grained and pebbly sandstone with angular to subrounded grains that are generally poorly sorted. The sandstone contains many interbeds and units of pebble-and-cobble conglomerate, dark-gray

siltstone, and mudstone intraclasts. Most of the conglomerate clasts are volcanic rock fragments, but some are granitic, and others are sandstone or limestone. Plant scraps and a few marine fossils have been found in the sandstone and its siltstone interbeds.

Downstream from the main sandstone outcrops on the Cheshnina River, and possibly overlying them, are green to gray lutites with thin sandstone interbeds at some places. These interbeds of sandstone are dark greenish gray and fine grained, and they locally have graded bedding and sole markings. Apparently the stratigraphically highest rocks are shaly and silty claystone and siltstone that contain brownish gray-weathering limestone interbeds and concretions.

The sandstone beds are estimated to be more than 1,000 feet, and are possibly more than 2,000 feet thick. Neither the top nor the base of the section is exposed. The lutites are at least several hundred feet thick, but in places they are in tight southeast-striking isoclinal and chevron folds, and their thickness could not be determined during the present reconnaissance.

These rocks were mapped as nonmarine and of Tertiary age by others (Moffit, 1939, pl. 2; Mendenhall, 1905, pl. 4). During the present studies, however, a belemnite and *Inoceramus* scraps were found in sandstone on the Chetaslina River, and several small pelecypods and small ammonites in outcrops on the Cheshnina River. Ralph W. Imlay, of the U.S. Geological Survey, states (written commun., 1963) that although the ammonites cannot be positively identified, they resemble immature forms of *Kepplerites*, or some related genus, of Callovian age, and that an early Late Jurassic age seems probable.

CRETACEOUS ROCKS

Clastic rocks of Early Cretaceous (Neocomian) age

Marine sedimentary rocks of Neocomian age form many small mountaintop outcrops between Long Glacier and Kuskulana Valley and constitute a larger area of continuous exposures near Kuskulana Pass (*fig. 2*). Correlative beds previously referred to as the tuffaceous units of Middle Jurassic age crop out south of the Chitina River in the northern Chugach Mountains. Moffit and Mertie (1923) described the Neocomian beds in some detail but were puzzled by their stratigraphic complexity and age and their relation to the Kotsina Conglomerate. Indeed, mollusks from these beds have been assigned by various paleontologists to the Middle Jurassic, Late Jurassic, Early Cretaceous, and Late Cretaceous (Moffit and Mertie, p. 44-48; Moffit, 1938, p. 66-70; Imlay and Reeside, 1954, p. 231).

The Neocomian beds are much less deformed than the latest Triassic and older rocks upon which they rest with angular unconformity, and near Kuskulana Pass (loc. E, fig. 2) they overlie granodiorite for which a potassium-argon age of 141 ± 5 m.y. was obtained on hornblende (sample 3, table 1). They are overlain, apparently unconformably, by beds of Albian age.

The Neocomian rocks are clastic marine deposits, but they are characterized by an almost white calcarenite facies which is composed of finely comminuted prisms of *Inoceramus* shell. Sandstone beds with lower proportions of biogenic calcite grains are brownish gray, greenish gray, or olive gray. North of the Chitina River these rocks are mostly fossiliferous, cross-bedded, calcareous sandstone and conglomerate. The thickest sections are from 600 to almost 1,000 feet thick. The sandstone and conglomerate intertongue greatly, and the stratigraphic sections differ widely from place to place (see Neocomian columnar sections, fig. 2). As with the nearby Kotsina Conglomerate, the conglomerate and sandstone clasts are chiefly greenstone, limestone, and argillite, and were derived from the subjacent formations. These rocks are nearshore, high-energy, shallow-water deposits.

North of the Chitina River, from 0 to 200 feet of rounded cobble-and-boulder conglomerate with a calcareous matrix, and in places containing many mollusks, occurs at the base of the Neocomian sequence. In the section north of the Kotsina River (loc. B, fig. 2) beds overlying the basal conglomerate are at least 600 feet thick. The lower 200–300 feet consist of calcareous sandstone that is fine to coarse grained and conglomeratic. The sandstone is fossiliferous, cross-bedded, and contains beds and lenses of calcarenite and of coquinoid sandstone. It in turn is overlain by a similar thickness of medium-dark and greenish-gray sandy siltstone with many, commonly large, mollusks. The upper part of the section consists of fossiliferous coarse sandstone, conglomerate, and calcarenite. At one place, about 60 feet of sandstone and conglomerate at the base of the upper coarse clastic rocks is overlain by about 50 feet of thick-bedded, very light gray (almost white), generally fine grained calcarenite with large *Inoceramus* shell fragments and some lenses of terrigenous pebbles and sand. The calcarenite section is overlain by a small thickness of incompletely exposed calcareous sandstone. Within one large outcrop the calcarenite interfingers with calcareous conglomerate, and it is evidently supplanted by fossiliferous calcareous sandstone and conglomerate within a quarter of a mile to the west. At one place the basal 5 feet of the calcarenite interfingers with pebble conglomerate and lenses out within a horizon-

tal distance of 6 to 8 feet. The calcarenite may represent a nearshore bar or bank of wave-comminuted shell fragments which abutted an area in which terrigenous detritus was being deposited in shallow water from a nearby stream mouth.

Correlative Neocomian beds in the northern Chugach Mountains south of the Chitina River and near its mouth (locs. C and D, fig. 2), are grossly similar in lithology to those north of the river. They are thicker, however, and include a thick unit of dark siltstone. Moffit (1914, p. 25-27) reported that these Neocomian rocks (loc. C, fig. 2) south of the Chitina River (which he considered Middle Jurassic) consisted of (1) a lower unit at least 500–600 feet thick composed of massive conglomerate with rounded pebbles and cobbles of argillite, diorite, greenstone, and quartz, set in a tuffaceous matrix, and (2) an upper unit several hundred feet thick of fossiliferous tuffaceous beds "... composed of dark fine-grained sandstone-like rock, slightly calcareous, and showing numerous small flakes of mica on the cleavage surface." Moffit mapped the upper unit as tuffaceous slate (1914, pl. 2), and tuffaceous shale (1938, pl. 2). He stated that the conglomerate appeared to rest unconformably upon Carboniferous(?) schist, but that the contact was poorly exposed and at many places appeared to be a fault. Moffit's description of these rocks suggests that they are calcareous conglomerate and siltstone lithologically like the Neocomian clastic rocks north of the Chitina River. However, three fossil collections from the upper unit were studied by T. W. Stanton (Moffit, 1914, p. 26), who referred them to the Middle Jurassic. These collections were reexamined during the present study and found to consist of fragmentary mollusks that could be either Jurassic or Cretaceous and that would not be out of place in the Hauterivian faunule collected from similar rocks about a mile to the east during the present study.

During the present reconnaissance the Neocomian rocks south of the Chitina River were examined only in a small stream at the east end of their outcrop area (loc. D, fig. 2) and their relationship to the rocks described by Moffit was not determined. Along this stream more than 800 feet of dark, fairly hard, fossiliferous glauconitic siltstone is exposed. It contains layers of thick *Inoceramus* shells and thin interbeds of calcareous sandstone and coarse siltstone crowded with *Inoceramus* prisms. This thick, fossiliferous siltstone probably was deposited somewhat farther from shore than the thinner and coarser equivalent rocks north of the river. The siltstone section is overlain by a few hundred feet of calcareous conglomerate and sandstone that contains abundant *Inoceramus* prisms

and has some interbeds of siltstone and of graywacke conglomerate and sandstone. As in the outcrops to the north, the conglomerate is dominated by clasts of altered volcanic rocks, limestone, black chert, and argillite, and contains some granitoid rocks and rarely quartz.

The abundant molluscan faunule in the lower and middle parts of the section (loc. B, fig. 2) north of the Kotsina River contains species of *Inoceramus*, *Pinna*, *Simbirskites*, and a belemnite which are probably of Hauterivian (late Neocomian) age. The same faunule was obtained from the dark siltstone south of the Chitina River (loc. D, fig. 2). The calcarenite near the top of the section north of the Kotsina River contains *Inoceramus* ovatoides of Anderson, which is of late Hauterivian to Barremian (late Neocomian) age. The genus *Simbirskites* in Europe is of middle to late Hauterivian age.

The Hauterivian beds of the southwestern Wrangell Mountains are correlative with the Nelchina Limestone and overlying dark sandstone in the Nelchina area of the Talkeetna Mountains, which Bergquist (1961) found to contain Hauterivian Foraminifera. The Nelchina Limestone is an almost white calcarenite characterized by *Inoceramus* shell fragments and prisms. The Nelchina Limestone overlies *Buchia*-bearing sandstone and conglomerate of Valanginian (early Neocomian) age which is absent in the Wrangell Mountains.

Siltstone, shale, and sandstone of Early Cretaceous (Albian) age

Albian-age rocks crop out in a broad syncline at Kuskulana Pass (loc. G, fig. 2), where they rest with apparent unconformity on the Neocomian beds just described, and in a small area on the southeast flank of Mount Drum (loc. A, fig. 2). The beds near Mount Drum have a moderate dip to the west and rest upon biotite granodiorite. Biotite from this pluton yielded a potassium-argon age of 126 ± 4 m.y. (sample 4, table 1), suggesting intrusion during Neocomian (Early Cretaceous) time. The fact that the unconformity at the base of the Hauterivian (upper Neocomian) beds in the Kotsina-Kuskulana Rivers area represents a greater structural discordance and a longer hiatus than the unconformity at the base of the overlying Albian beds suggests that the pluton was intruded before deposition of the upper Neocomian beds of the Kotsina-Kuskulana Rivers area, and that it was unroofed by a combination of pre-Hauterivian and pre-Albian erosion.

The Albian rocks at Kuskulana Pass are similar to strata cropping out near Fourth of July Pass in the upper Chitina Valley that have been designated the

Kennicott Formation. However, usage of the name 'Kennicott has been so varied that use of the term should be suspended until detailed mapping is accomplished in the type area. At Kuskulana Pass the Albian rocks (loc. G, fig. 2) are about 1,500 feet thick and are dominantly siltstone. Some sandstone may occur at the base of the section, but the lowest beds are poorly exposed at the place visited. The bulk of the section is siltstone, silty claystone, and shale that are slightly greenish gray and medium dark gray and weather to medium gray, brownish gray, or greenish gray. The siltstone contains limestone concretions and lentils and a few volcanic ash layers. An interval 150 feet thick in the lower part of the sequence contains abundant shells of the pelecypod *Arcellina* in limestone concretions and coquinoid beds. Another interval at least 200 feet thick in the middle of the sequence contains 5 to 10 percent of thin to thick interbeds of coarse siltstone, sandstone, and minor conglomerate. These interbeds resemble turbidites, being commonly graded, and many show sole markings, wavy bedding, small-scale crossbedding, and channeled bases. The uppermost exposed beds in this section are siltstone and silty claystone with limestone concretions, but the top of the section is not preserved. Fossils were not obtained from the highest beds, and thus their age is unknown. A few observations along the northeast side of the syncline at Kuskulana Pass suggest that sandstone and conglomerate interbeds in the middle of the sequence may increase in number and thickness to the north.

The Albian strata near Mount Drum are lithologically quite different from those at Kuskulana Pass, for they are characterized by very carbonaceous silt and clay shales and contain beds of granite wash. They are also slightly younger. The exposed section (loc. A, fig. 2) exceeds 650 feet in thickness. Its basal part is 350 feet thick and consists mostly of gray, soft siltstone and silt shale resting on granodiorite. It contains large limestone concretions with marine fossils, wood fragments, and glauconite grains. Medium to very thick interbeds of arkose, mostly coarse and pebbly, and of granite wash are distinctive features of this unit. The thickest of these interbeds are composed of angular and subangular granitic blocks mixed with intraclasts of sandstone and limestone, yet they are separated from the underlying granodiorite by tens of feet of shallow marine siltstone. These relationships suggest that the basal Albian sea floor had appreciable relief, and that the coarse detritus was dumped in and buried without significant reworking, even though the beds are apparently of shallow marine origin.

Dark, very carbonaceous, and commonly fissile brownish-gray-weathering shale and siltstone, 175 feet thick, overlie the basal unit. These rocks have yielded some animal trails but no other fossils. The carbonaceous character of these beds, the presence of trails, and the absence of mollusks within them suggest that they are of brackish-water or lagoonal origin. The uppermost 140 feet of the exposed section is dark, chunky, and platy weathering siltstone that contains large limestone concretions, some of which contain marine fossils. The top of the unit is unexposed, but the section is overlain by Cenozoic volcanic rocks.

The basal beds near Mount Drum contain *Breweriaceras hulenense* and other ammonites of the *Breweriaceras hulenense* local faunizone of late early Albian age. The upper beds contain *Puzosia alaskana* Imlay of probably the same age. *Aucellina*, which is restricted to the lower lower Albian zone of *Moffittites robustus*, is apparently absent from the Mount Drum section, which is therefore slightly younger than the *Aucellina*-bearing Albian beds at Kuskulana Pass. The Mount Drum section is also lithologically different from that at Kuskulana Pass; it more closely resembles the Albian beds of the northern part of the Nelchina area, which consist of coaly beds overlain by very fossiliferous shallow marine shale. The lithologic differences between the Albian rocks of Kuskulana Pass and Mount Drum suggest that the former should be named according to the formal stratigraphic nomenclature applied to the upper Chitina Valley, and the latter according to the nomenclature of the Nelchina area and Copper River lowland.

The upper Mesozoic rocks of the southwestern Wrangell Mountains record at least three episodes of deep erosion followed by sedimentation. These episodes are recorded at (1) the base of the Kotsina Conglomerate, (2) the base of the Neocomian beds, (3) the base of the Albian beds. These rocks also record evidence of plutonism of latest Early or earliest Middle Jurassic age, of Late Jurassic age, and of Early Cretaceous age within or near the Matanuska geosyncline. These rocks have some overall lithologic similarities to rocks of similar age in nearby areas,

but their stratigraphic sequence is nevertheless quite different from that found in the nearby areas. These relations illustrate that these rocks were deposited in a tectonically active trough that typically produced rocks that are lithologically similar over wide areas, but which are arranged in highly distinctive and variable local sequences. Long-range extrapolation of local stratigraphic details in these rocks is, therefore, a risky enterprise.

REFERENCES

- Bergquist, H. R., 1961, Early Cretaceous (middle Neocomian) microfossils in south-central Alaska: Art. 374 in U.S. Geol. Survey Prof. Paper 424-D, p. D236-D237.
- Detterman, R. L., Reed, B. L., and Lanphere, M. A., 1965, Jurassic plutonism in the Cook Inlet region, Alaska, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-D, p. D16-D21.
- Grantz, Arthur, Jones, D. L., and Lanphere, M. A., 1965, Upper Mesozoic rocks of the lower Chitina Valley, Alaska [abs]: Geol. Soc. America Spec. Paper 82, p. 255 and 356.
- Grantz, Arthur, Thomas, Herman, Stern, T. W., and Sheffey, N. B., 1963, Potassium-argon and lead-alpha ages for stratigraphically bracketed plutonic rocks in the Talkeetna Mountains, Alaska: Art. 16 in U.S. Geol. Survey Prof. Paper 475-B, p. B56-B59.
- Imlay, R. W., 1952, Correlation of the Jurassic Formations of North America, exclusive of Canada: Geol. Soc. America Bull., v. 63, p. 953-992.
- Imlay, R. W., and Reeside, J. B., Jr., 1954, Correlation of the Cretaceous formations of Greenland and Alaska: Geol. Soc. America Bull., v. 65, p. 223-246.
- Mendenhall, W. C., 1905, Geology of the central Copper River region, Alaska: U.S. Geol. Survey Prof. Paper 41, 133 p.
- Moffit, F. H., 1914, Geology of the Hanagita-Bremner region, Alaska: U.S. Geol. Survey Bull. 576, 56 p.
- Moffit, F. H., 1938, Geology of the Chitina Valley and adjacent area, Alaska: U.S. Geol. Survey Bull. 894, 137 p. [1939].
- Moffit, F. H., and Maddren, A. G., 1909, Mineral resources of the Kotsina-Chitina region, Alaska: U.S. Geol. Survey Bull. 374, 103 p.
- Moffit, F. H., and Mertie, J. B., Jr., 1923, The Kotsina-Kuskulana district, Alaska: U.S. Geol. Survey Bull. 745, 149 p.
- Rohn, Oscar, 1900, A reconnaissance of the Chitina River and the Skolai Mountains, Alaska: U.S. Geol. Survey 21st Ann. Rept., pt. 2, sect. h, p. 393-440, pls. 51-54.
- Schrader, F. C., and Spencer, A. C., 1901, The geology and mineral resources of a portion of the Copper River district, Alaska: U.S. Geol. Survey, spec. pub., 94 p.

